

Sri Hidayati¹, Subeki Subeki¹, M. Perdiansyah Mulia Harahap², Sutopo Hadi³

Utilization of Lignin Monomer Isolation from Black Liquor of Empty Palm Oil Bunches as a Prebiotic

Upotreba izoliranog monomera lignina iz crnog luga dobivenoga od praznih grozdova palmina ploda kao prebiotika

ORIGINAL SCIENTIFIC PAPER

Izvorni znanstveni rad

Received – prispjelo: 17. 2. 2022.

Accepted – prihvaćeno: 20. 4. 2022.

UDK: 630*86; 665.947.4

<https://doi.org/10.5552/drvind.2023.0015>

© 2023 by the author(s).

Licensee Faculty of Forestry and Wood Technology, University of Zagreb.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

ABSTRACT • Waste is generated in the form of black liquor during pulp processing. Furthermore, lignin, the main ingredient of black liquor, contains a phenylpropanoid compound with prebiotic and antimicrobial activity. Also, it is a component of lignocellulose that has prebiotic and antimicrobial activity effects due to its indigestible nature and it comprises phenylpropanoid components. Therefore, this study examines and identifies the lignin purification and test results of its prebiotic activity. The technique used to identify lignin fraction is called Thin Layer Chromatography (TLC) and Gas Chromatography-Mass Spectrometry (GC-MS) plates. Prebiotic activity tests were performed using the calculation of total bacteria on the growth of *Lactobacillus casei*. The results showed that the purification process using CHCl_3 , 3 % MeOH: CHCl_3 , 20 % MeOH: CHCl_3 , and MeOH yielded 10.68 %, 6.34 %, 11.38 % 44.85 %, respectively. The 3 % MeOH: CHCl_3 fraction contained benzaldehyde, 4-hydroxy-3,5-dimethoxy, 1-methylbutyl hexadecanoate, oleic acid, and di-2-ethylhexyl phthalate. The 3 % MeOH: CHCl_3 fractionate with a concentration of 15 % also showed a prebiotic activity for *L. casei* at 6.1×10^2 colonies/mL.

KEYWORDS: black liquor; lignin monomer; empty palm oil bunch; prebiotic

SAŽETAK • Tijekom prerade pulpe nastaje otpad u obliku crnog luga. Lignin kao glavni sastojak crnog luga sadržava spoj fenilpropanoid prebiotičkoga i antimikrobnog djelovanja. Također, on je i dio lignoceluloze koja zbog svoje neprobavljivosti ima prebiotičko i antimikrobno djelovanje, a sastoji se od fenilpropanoidnih komponentata. Stoga se u ovom istraživanju ispituju i razmatraju pročišćivanje lignina i rezultati prebiotičke aktivnosti. Za identifikaciju frakcija lignina primijenjene su tankoslojna kromatografija (TLC) i plinska kromatografija s masenom spektrometrijom (GC-MS). Ispitivanje prebiotičke aktivnosti provedeno je korištenjem izračuna ukupnog broja bakterija na rast *Lactobacillus casei*. Rezultati su pokazali da je proces pročišćivanja uz pomoć CHCl_3 , 3 % MeOH: CHCl_3 , 20 % MeOH: CHCl_3 , i MeOH dao prinosa od 10,68 %, 6,34 %, 11,38 % i 44,85 %. Frakcija 3 %

¹ Authors are researchers at Lampung University, Agricultural Technology Department, Bandar Lampung, Indonesia. <https://orcid.org/0000-0002-8790-4322>, <https://orcid.org/0000-0002-3938-2945>

² Author is researcher at Lampung State Polytechnic, Bandar Lampung, Indonesia. <https://orcid.org/0000-0002-7384-236X>

³ Author is researcher at Lampung University, Department of Chemistry, Bandar Lampung, Indonesia. <https://orcid.org/0000-0001-6464-7215>

MeOH:CHCl₃ sadržavala je benzaldehid, 4- hidroksi-3,5-dimetoksi, 1-metilbutil heksadekanoat, oleinsku kiselinu i di-2-etilheksil ftalat. Frakcija 3 % MeOH:CHCl₃ koncentracije 15 % također je pokazala prebiotičku aktivnost za *L. casei* pri $6,1 \times 10^3$ kolonija/mL.

KLJUČNE RIJEČI: crni lug; monomer lignina; prazni grozdovi palmina ploda; prebiotik

1 INTRODUCTION

1. UVOD

During pulp processing, liquid waste is generated in black liquor, which has lignin content with a complex structure that is difficult to decompose; thus, it can pollute the environment (Lara *et al.*, 2003). However, the lignin content is approximately 25-35 % of the total black liquor (Lara *et al.*, 2003; Goujon *et al.*, 2003). This implies that it can be isolated for further use. Currently, people want to use black liquor as a source of raw material for lignin (Min *et al.*, 2013), which is further used as biomass (Schorr *et al.*, 2014), adhesives (Ramires *et al.*, 2010; Uraki and Koda, 2015), dispersants, surfactants, antioxidants in plastics and rubbers, dyes, synthetic floors, thermosets, paints, and fuel for road maintenance (Laurichesse and Averous, 2013; Mankar *et al.*, 2012; Magnus and Hakan, 2014; Padkościelna *et al.*, 2017; Baurhoo *et al.*, 2008).

Hidayati *et al.* (2018) isolate lignin from black liquor of empty palm oil bunches by using NaOH at a concentration of 30 % yielded of 5.67 %, with a pH of 5.42, total solid content of black liquor 65.11 %, methyl lignin level of 14.61 %, and equivalent weight 1787.23. Prayuwidayati *et al.* (2016) showed that lignin is used as a prebiotic, which is considered to have an antimicrobial effect by reducing the growth rate of pathogenic bacteria in the cattle rumen. According to Baurhoo *et al.* (2008), the tested, purified lignin product has a prebiotic effect. Furthermore, prebiotic is a beneficial component of human food because it enhances the growth and activity of several bacteria in the colon, thus improving the health of the human digestive tract (Toma and Pokrotnieks, 2006). This study aims to identify the components contained in lignin purification products by TLC, Chromatography Column, and GC-MS as testing the prebiotic activity.

2 MATERIALS AND METHODS

2. MATERIJALI I METODE

2.1 Tools and materials

2.1.1. Oprema i materijali

The materials used were the Empty Palm Oil Bunches (EPOB) black liquor resulting from formacell pulping, acetic acid, formic acid, HCl, CuSO₄, H₂O₂, NaOH, pyridine, MeOH, CHCl₃, and equates, as well as NA (Nutrient Agar), NB (Nutrient Broth), media de Man, Rogosa and Sharpe Agar (MRSA), and media de Man, Rogosa and Sharpe Broth

(MRSB). Black liquor was obtained from the formacell pulping process using the method used by Hidayati *et al.* (2017). The advantages of formacell pulping compared to other pulping methods are that it is more environmentally friendly, the pulp yield is high, the pulp produced is sulfur free, and it has high brightness and good strength. The pulping process was carried out in a 1.3 liter autoclave with 20 g of EPOB fiber and liquid for a sample ratio of 15:1 flat bottom, and a wide-mouth boiling flask equipped with a condenser. The solvent ratio of acetic acid: formic acid (85:15) and HCl catalyst was given 0.5 % for 1 hour of cooking time at 130 °C. After cooking, the remaining boiled liquid was separated and collected by filtering. The microbial cultures used were *L. casei*. Furthermore, the tools used were oven, Buchner funnel, silica gel 60 F₂₅₄ TLC plate, column chromatography, chamber, autoclave, micropipette, incubator, calipers, capillary pipette, GCMS (Varian/CP- 3800 GC and Saturn 2200 MS), and supporting glasses.

2.2 Study method

2.2.1. Ispitne metode

The study was conducted with a Completely Randomized Design (CRD) in which black liquor was extracted until lignin was obtained twice. Subsequently, lignin was fractionated until a monomer fraction was obtained. TLC analysis identified each lignin monomer fraction, and the prebiotic activity was screened for *L. casei*. The fraction with the highest prebiotic activity was analyzed using GC-MS to determine the chemical composition. Furthermore, they were generated at several concentrations of 0 %, 2.5 %, 5 %, 7.5 %, 10 %, 12.5 %, and 15 %, and they were to be tested for antimicrobial and prebiotic activity for three replications. The data obtained were analyzed descriptively and presented in figures and tables.

2.3 Study implementation

2.3.1. Provedba ispitivanja

2.3.1.1 Lignin degradation

2.3.1.1.1. Degradacija lignina

The lignin obtained from black liquor due to the pulp cooking with the raw material of EPOB was precipitated and degraded using CuSO₄, pyridine, and H₂O₂. A total of 2 g of lignin precipitate was dissolved with 20 mL NaOH solution to a pH of 12. Afterward, a total of 25 mL of 10⁻² M CuSO₄ and 5 mL pyridine were added to the lignin solution, and then it was stirred with a magnetic stirrer for 30 minutes. 10 ml of




Figure 1 Lignin monomer fractionation flow diagram
Slika 1. Dijagram toka frakcioniranja monomera lignina

H₂O₂ 1 M was also added five times for 30 minutes, stirred, and stored in an unlighted room for 72 hours.

2.3.2 Lignin purification fraction

2.3.2. Pročišćivanje frakcija lignina

The fractionated filtrate was evaporated until a residue was obtained and dissolved in 500 mL H₂O. After that, extraction was performed with EtOAc of 500 mL for three times to obtain H₂O and EtOAc layers that were evaporated until a residue was obtained. The residue was then put into the chromatographic column silica gel and eluted with CHCl₃ (1L) solution. Furthermore, the CHCl₃ fraction was then evaporated, and the obtained residue was purified by preparative thin-layer chromatography (PTLC). The lignin monomer isolation process can be seen in Figure 1.

2.3.3 Lignin fraction identification

2.3.3. Identifikacija frakcija lignina

Identification of lignin monomers obtained was made using TLC and GC-MS plates. Subsequently, the isolates were dissolved in hexane and injected into GC-MS devices. The analysis was performed using GC-MS CP-3800 variant with MS Saturn 2200 detector, and VF-5 ms column 30 mm × 0.25 mm by manual injection method at 240 °C for 40 minutes (Suroso *et al.*, 2018).

2.4 Prebiotic activity test for lignin fraction

2.4. Ispitivanje prebiotičke aktivnosti frakcija lignina

Prebiotic tests were performed microbiologically using *L. casei* grown on MRSA media, and the meas-

ured variable was the number of bacteria counted using the living bacterial colonies method. The lignin monomer fraction was composed of media such that each cup was 0 %, 2.5 %, 5 %, 7.5 %, 10 %, 12.5 %, and 15 %. *L. casei* that had been refreshed on MRSB media was diluted as much as 0.05 mL into 4.95 mL of physiological solution. Then, 1 mL of physiological solution was inoculated into MRSA media, and it was poured into 20 petri cups of 19.5, 19, 18.5, 18, 17.5, and 17 mL and 3 % MeOH:CHCl₃ fraction of 0, 0.5, 1, 1.5, 2, 2.5, and 3 mL were added. Furthermore, the bacteria were incubated for 24 hours in an incubator. The number of bacteria was calculated by counting the living bacterial colonies using the following method of Ogimoto and Imai (1981).

$$\text{Microbe population} = \frac{\text{Colony account (kol)}}{0.05 \times 10^x \times 0.1 \text{ (ml)}} \quad (1)$$

Note: x: tube xth retail series

3 RESULTS AND DISCUSSION

3. REZULTATI I RASPRAVA

3.1 Lignin isolation

3.1. Izoliranje lignina

The black liquor produced by formacell pulping was then isolated to obtain lignin isolates. Furthermore, the extraction process of lignin from black liquor was performed by adding water to 500 mL of the sample in a 2 L glass beaker and allowed to form a precipitate which was then separated by filtering and dried for three days at room temperature; afterwards, it was observed for its characteristics (Table 1).

Table 1 Characteristics of isolated lignin**Tablica 1.** Svojstva izoliranog lignina

Parameter / Parametar	Characteristic / Svojstvo
Color / boja	Black / crna
Form / konzistencija	Solid / čvrsta
pH (25 °C)	4.5
Yield / prinosa	1.74%
Water content / sadržaj vode	0.24%

Lignin isolates obtained have a black appearance, and they are solids in the form of fiber crumbs. They have a moisture content of 0.24 %. The yield of lignin obtained is 1.74 %, and this is presumed because there was no addition of acid to the precipitation process.

3.2 Lignin monomer fraction screening for *L. casei*

3.2. Probir frakcije monomera lignina za *L. casei*

The process of lignin degradation was performed by dissolving it into a solution of CuSO_4 , pyridine, and NaOH, after which it was stirred and H_2O_2 was added. The obtained fraction was then added to the chromatographic column silica and eluted with CHCl_3 , 3 % MeOH: CHCl_3 , 20 % MeOH: CHCl_3 , and MeOH, respectively. Then, each fraction was dried. The fraction of CHCl_3 , 3 % MeOH: CHCl_3 , 20% MeOH: CHCl_3 , and MeOH yielded 10.68 %, 6.34 %, 11.38 %, and 44.85 %, respectively. The results of each fraction were identified on a TLC plate using UV light to examine the content of chemical compounds qualitatively.

Black spots on the TLC showed the presence of chemical compounds. Furthermore, the spot on TLC of each fraction consists of different compounds based on their retention time. For example, the spots generated at fractions of CHCl_3 , 3 % MeOH: CHCl_3 , 20 % MeOH: CHCl_3 , and MeOH sequentially produced 4, 4, 2, and 1 spot, which was considered to be the dominant compound in each lignin fraction. Figure 2 and Table 2 show the black spot on TLC and the screening results for *L. casei* growth, respectively (Table 2).




Figure 2 Chromatographic profile of thin layers of each fraction: a) CHCl_3 ; b) 3 % MeOH: CHCl_3 ; c) 20 % MeOH: CHCl_3 ; d) MeOH

Slika 2. Kromatografski profil tankih slojeva pojedine frakcije: a) CHCl_3 ; b) 3 % MeOH: CHCl_3 ; c) 20 % MeOH: CHCl_3 ; d) MeOH

Table 2 Screening lignin fraction as a prebiotic for *L. casei***Tablica 2.** Probir frakcije monomera lignina za *L. casei*

Fraction / Frakcija	Number of microbes, 10^2 colony/mL / Broj mikroba, 10^2 kolonija/mL
CHCl_3	1.48 ± 0.29
3 % MeOH: CHCl_3	4.52 ± 0.10
20 % MeOH: CHCl_3	2.58 ± 0.23
MeOH	2.41 ± 0.34

The 3 % MeOH: CHCl_3 fraction showed higher prebiotic activity than other fractions. It showed prebiotic activity against *L. casei* at 4.52×10^2 colonies/mL, while the CHCl_3 fraction, 20% MeOH: CHCl_3 , and MeOH showed prebiotic activity against *L. casei* at 1.48×10^2 colonies/mL, 2.58×10^2 colonies/mL, and 2.41×10^2 colonies/mL, respectively.

3.3 Identification of 3 % MeOH: CHCl_3 fraction compound content

3.3. Identifikacija spojeva u frakciji 3 % MeOH: CHCl_3

The identification process was performed by injecting a 3 % MeOH: CHCl_3 fraction into the GC-MS (Figure 3).

The 3 % MeOH: CHCl_3 fraction contains (a) benzaldehyde, 4-hydroxy-3,5-dimethoxy- 2.76 %, (b) 1-methylbutyl hexadecanoate as much as 41.03 %, (c) oleic acid as much as 3.61 %, and (d) di-2-ethylhexyl phthalate as much as 31.25 %. Furthermore, the main product of the hydrolysis of lignin using NaOH catalysts is monomeric phenols and oligomers (Roberts *et al.*, 2011). Figure 4 and Table 3 show the compound structure of the 3 % MeOH: CHCl_3 fraction and the identification results, respectively.

3.4 Prebiotic activity of 3 % MeOH: CHCl_3 fraction against *L. casei*

3.4. Prebiotičko djelovanje frakcije 3 % MeOH: CHCl_3 na *L. casei*

Prebiotic activity test uses a concentration between growth media with a fraction of 3% methanol-chloroform of 0, 2.5, 5, 7.5, 10, 12.5, and 15 (%). The




Figure 3 Fraction chromatogram of 3 % MeOH:CHCl₃: a) benzaldehyde, 4-hydroxy-3,5-dimethoxy; b) 1-methylbutyl hexadecanoate; c) oleic acid; d) di-2-ethylhexyl phthalate

Slika 3. Kromatogram frakcije 3 % MeOH:CHCl₃: a) benzaldehid, 4-hidroksi-3,5-dimetoksi; b) 1-metilbutil heksadekanoat; c) oleinska kiselina; d) di-2-etilheksil ftalat




Figure 4 Fraction compound of 3% MeOH:CHCl₃ includes: a) benzaldehyde, 4-hydroxy-3,5-dimethoxy; b) 1-methylbutyl hexadecanoate; c) oleic acid and d) di-2-ethylhexyl phthalate

Slika 4. Spojevi u frakciji 3 % MeOH:CHCl₃ uključuju: a) benzaldehid, 4- hidroksi-3,5-dimetoksi; b) 1-metilbutil heksadekanoat; c) oleinsku kiselinu i d) di-2-etilheksil ftalat

number of colonies from the growth of *L. casei* for each concentration of 3 % MeOH:CHCl₃ fraction is shown in Figure 5.

The 3 % MeOH:CHCl₃ fraction at a concentration of 15 % showed the highest prebiotic activity of 6.10×10^2 colonies/mL against *L. casei*. In comparison, the 3 % MeOH:CHCl₃ fraction at a concentration of 2.5 % showed the lowest prebiotic activity of $2,69 \times 10^2$ colonies/mL against *L. casei*. The prebiotic activity of the 3 % MeOH:CHCl₃ fraction on the growth of *L. casei* is shown in Figure 6.

The results of the analysis of the prebiotic activity of the 3 % MeOH:CHCl₃ fraction showed that the

higher the concentration of the growth medium, the higher the growth of *L. casei*. Baurhoo *et al.* (2008) stated that the lignin monomer in the alcell process acts as a prebiotic. Prayuwidayati *et al.* (2016) reported that formacell purification products are generated from EPOB and oligomers, which are phenolic compounds that stimulate the growth of lactic acid bacteria. This shows that the isolated fraction of lignin and its derivatives can replace glucose in MRSA medium, and it is used as an energy source for the growth of *L. casei*. These also showed that the isolated lignin fraction could be used as a feed supplement and support the growth of *L. casei*. Lignin in the purified alcell

Table 3 Identification of lignin fraction monomer compounds of 3 % MeOH:CHCl₃**Tablica 3.** Identifikacija spojeva u frakciji 3 % MeOH:CHCl₃ monomera lignina

Retention time <i>Vrijeme retencije</i>	Molecule weight <i>Molekulska masa</i>	Compound / Spoj	%
8.886	207	Phenol.2-(1-methylpropyl)-.methylcarbamate	0.12
11.008	212	Propanoic acid.3-chloro-.4-formylphenyl ester	0.38
11.942	152	Vanillin	1.96
13.958	166	Ethanone.1-(4-hydroxy-3-methoxyphenyl)-	0.15
14.709	166	Undecanoic acid.10-methyl-.methyl ester	0.47
15.839	214	Benzoic acid.4-hydroxy-3-methoxy-	1.04
16.313	168	Diethyl phthalate	0.13
17.952	222	Benzaldehyde.4-hydroxy-3.5-dimethoxy-	2.76
18.621	182	p-Anisic acid.4-nitrophenyl ester	0.32
18.838	273	m-Anisic acid.3.4-dichlorophenyl ester	0.90
19.022	413	Carbamic acid.N-[1.1- bis(trifluoromethyl)ethyl]-4-(1.1.3.3-tetra-methylbutyl)phenyl ester	0.95
19.114	220	4-Methyl-2-tert-octylphenol	0.54
19.235	296	m-Anisic acid.3.4-dichlorophenyl ester	1.10
19.447	270	Hexestrol	0.58
19.643	220	Phenol.2-methyl-4-(1.1.3.3- tetramethylbutyl)-	0.33
19.708	192	1.3-Dimethyl-5-ethyladamantane	0.74
19.870	413	Carbamic acid.N-[1.1- Bis (trifluoromethyl)ethyl]-4-(1.1.3.3- tetra-methylbutyl)phenyl ester	1.06
20.034	220	Phenol.2-methyl-4-(1.1.3.3- tetramethylbutyl)-	0.71
20.126	228	Tetradecanoic acid	1.17
21.690	268	2-Pentadecanone.6.10.14-trimethyl-	0.07
21.980	194	Caffeine	0.12
22.135	278	1.2-Benzenedicarboxylic acid.bis(2- methylpropyl) ester	0.70
22.520	338	Erucic acid	0.09
22.848	604	Tritetracontane	0.28
22.968	268	9-Hexadecenoic acid. methyl ester.(Z)-	0.41
23.116	276	7.9-Di-tert-butyl-1-oxaspiro(4.5)deca-6.9-diene-2.8 dione	1.60
23.395	270	Pentadecanoic acid.14-methyl-.methyl ester	1.37
23.358	292	Benzenepropanoic acid.3.5-bis(1.1-dimethylethyl)-4 hydroxy .methyl ester	0.20
24.659	326	1-Methylbutyl hexadecanoate	41.03
26.713	298	1-Eicosanol	0.23
26.837	294	9.12-Octadecadienoic acid (Z,Z)-.methyl ester	0.32
26.971	352	9.12.15-Octadecatrienoic acid.2.3 dihydroxypropylester.(Z.Z.Z)-	0.58
28.036	282	Oleic acid	0.89
28.453	282	Oleic acid	3.61
28.853	282	Oleic acid	0.33
30.953	604	Tritetracontane	0.21
31.997	324	4.8.12.16-tetramethylheptadecan-4-olide	0.57
32.860	298	1-Eicosanol	0.11
32.963	604	Tritetracontane	0.16
35.399	242	1-decanol.2-hexyl-	0.25
36.624	390	Di-2-ethylhexyl phthalate	31.25
38.491	592	1-Hentetracontanol	0.21

process showed a prebiotic effect on chickens, promoting the growth of beneficial bacteria and improving intestinal morphological structure, as measured by an increase in the number of villi and goblet cells (Baurhoo *et al.*, 2008). Cotta and Whitefield (1998) stated that all hemicellulolytic rumen bacteria could utilize xylooligosaccharides as growth substrates. Therefore, increased digestibility of crude protein and fiber supports ruminant protein and energy metabo-

lism. The non-digestible carbohydrate prebiotics includes lactulose, inulin, resistant starch, and a number of oligosaccharides, a source of carbohydrates for beneficial bacteria in the digestive tract (Crittenden, 1999). Substrates such as inulin, fructooligosaccharides (FOS), and mannan-oligosaccharides (MOS), derived from yeast cells, absorbed the host and were hydrolyzed by endogenous digestive enzymes. The possible mechanism is a decrease in pH due to the




Figure 5 Effect of several concentrations of 3 % MeOH:CHCl₃ fraction as a prebiotic on the number of *L. casei* microbes
Slika 5. Utjecaj nekoliko koncentracija frakcije 3 % MeOH:CHCl₃ na broj mikroba *L. casei*




Figure 6 Prebiotic activity of 3 % MeOH:CHCl₃ fraction on the growth of *L. casei*
Slika 6. Utjecaj prebiotičke aktivnosti frakcije 3 % MeOH:CHCl₃ na rast *L. casei*

production of short-chain fatty acids, bacteriocin secretion, and immune stimulation.

Meanwhile, MOS as a prebiotic has a different mechanism that does not cause an increase in the population of beneficial bacteria, due to its ability to attach to mannose-specific lectins of G-negative type 1 fimbriae pathogens such as *Salmonella* and *E. coli*, which is excreted from the digestive tract (Toma and Prokotiņeks, 2006; Crittenden, 1999). Similar to the effect of MOS and antibiotic-free diet supplemented with 1.25 % alcell lignin (LL) on *Lactobacilli* and *Bifidobacteria*, lignin at low levels is potentially classified as a prebiotic (Baurhoo *et al.*, 2008). The results showed that using lignin in the alcell process (12.5 g) increases the concentration of intestinal *Lactobacilli* and *Bifidobacteria* in broilers (Baurhoo *et al.*, 2008). In addition, the proliferation of gastrointestinal microbiota populations

such as *Lactobacilli* and *Bifidobacteria* was increased through the consumption of prebiotics. The increase in the number of microbiotic in the feces ranges from 10-100 times (Thomas *et al.*, 2004).

4 CONCLUSIONS

4. ZAKLJUČAK

Lignin, the main ingredient of black liquor generated from bunches of empty oil palm fruit, contains benzaldehyde, 4-hydroxy-3,5-dimethoxy, 1-methylbutyl hexadecanoate, oleic acid, and di-2-ethylhexyl phthalate. Furthermore, its fraction of 3 % MeOH:CHCl₃ with a concentration of 15 % showed prebiotic activity against *L. casei* of 6.1×10^2 colonies/m. These results indicated that the lignin monomer is potentially prebiotic.

Acknowledgements – Zahvala

Thank you to the Ministry of Education and Culture of the Republic of Indonesia for the financial assistance, the laboratory for testing the quality of agricultural products, the Faculty of Agriculture for the support of laboratory facilities.

5 REFERENCES

5. LITERATURA

- Baurhoo, B.; Ruiz-Feria, C. A.; Zhao, X., 2008: Purified lignin: Nutritional and health impacts on farm animals – A review. *Animal Feed Science and Technology*, 144: 175-184. <https://doi.org/10.1016/j.anifeedsci.2007.10.016>
- Crittenden, R. G., 1999: Prebiotics. Tannock, G. W. (ed): *Probiotic: A Critical Review*. Horizon Scientific Press, Wymondham, pp. 141-156.
- Cotta, M. A.; Whitefield, T. R., 1998: Xylooligosaccharides utilization by ruminal anaerobic bacterium *Selemonas ruminantium*. *Current Microbiology*, 36: 183-189. <https://doi.org/10.1007/s002849900291>
- Ramires, E. C.; Megiatto Jr., J. D.; Gardrat, C; Castellán, A.; Frollini, E., 2010: Valorization of an industrial organosolv-sugarcane bagasse lignin: Characterization and use as a matrix in biobased composites reinforced with sisal fibers. *Biotechnology and Bioengineering*, 107 (4): 612-621. <https://doi.org/10.1002/bit.22847>
- Goujon, T.; Ferret, V.; Mila, I.; Pollet, B.; Ruel, K.; Burlat, V.; Joseleau, J. P.; Barrière, Y.; Lapiere, C.; Jouanin, L., 2003: Down-regulation of the AtCCR1 gene in *Arabidopsis thaliana*: effects on phenotype, lignins and cell wall degradability. *Planta*, 217: 218-228. <https://doi.org/10.1007/s00425-010-1202-1>
- Hidayati, S.; Zuidar, A. S.; Satyajaya, W.; Murhadi, Retnowati, D., 2018: Isolation and characterization of formacell lignins from oil empty fruits bunches. *IOP Conference Series: Materials Science and Engineering*, 344: 012006 <https://doi.org/10.1088/1757-899X/344/1/012006>.
- Hidayati, S.; Zuidar, A. S.; Satyajaya, W., 2017: Effect of acetic acid: formic acid ratio on characteristics of pulp from oil palm empty fruit bunches (OPEFB). *Journal of Engineering and Applied Science*, 12 (2): 3802-3807.
- Lara, M. A.; Rodríguez-Malaver, A. J.; Rojas, O. J.; Holmuist, O.; González, A. M.; Bullón, J., 2003: Blackliquor lignin biodegradation by *trametes elegans*. *International Biodeterioration and Biodegradation*, 52: 167-173.
- Laurichesse, S.; Averous, L., 2013. Chemical modification of lignin: Toward Biobased Polymers. *Progress in Polymer Science*, 39 (7): 1266-1290. <http://dx.doi.org/10.1016/j.progpolymsci.2013.11.004>
- Mankar, S. S.; Chaudhari, A. R.; Soni, I., 2012: Lignin in phenolformaldehyde adhesives. *International Journal of Knowledge Engineering*, 3 (1): 116-118.
- Magnus, N.; Hakan, E., 2014: Lignin: recent advances and emerging applications. *Current Opinion in Colloid & Interface Science*, 19 (5): 409-416. <https://doi.org/10.1016/j.cocis.2014.08.004>
- Min, D. Y.; Smith, S. W.; Chang, H. M.; Jameel, H., 2013: Influence of isolation condition on structure of milled wood lignin characterized by quantitative C nuclear magnetic resonance spectroscopy. *BioResources*, 8 (2): 1790-1800.
- Ogimoto, K.; Imai, S., 1981: *Atlas of rumen microbiology*. Japan Scientific Societies Press, Tokyo, pp. 231.
- Podkościelna, B.; Goliszek, M.; Sevastyanova, O., 2017: New approach in the application of lignin for the synthesis of hybrid materials. *Pure and Applied Chemistry*, 89 (1): 161-171. <https://doi.org/10.1515/pac-2016-1009>
- Prayuwidayati, M.; Sunarti, T. C.; Sumardi, Subeki, Wiryawan, K. G., 2016: Use of lignin formacell of empty bunch palm fiber as feed supplement and prebiotics Candidate in Ruminant. *Pakistan Journal of Nutrition*, 15 (1): 58-65. <https://doi.org/10.3923/pjn.2016.58.65>
- Roberts, V.; Stein, V.; Reiner, T.; Lemonidou, A.; Li, X.; Lercher, J. A., 2011: Towards quantitative catalytic lignin depolymerization. *Chemistry: A European Journal*, 17 (21): 5939-5948. <https://doi.org/10.1002/chem.201002438>
- Schorr, D.; Diouf, P. N.; Stevanovic, T., 2014: Evaluation of industrial lignins for biocomposites production. *Industrial Crops and Products*, 52: 65-73. <https://doi.org/10.1016/j.indcrop.2013.10.014>
- Suroso, E.; Utomo, T. P.; Hidayati, S.; Nuraini, A., 2018: Smoked mackerel using liquid smoke from red-distilled rubber wood. *Jurnal Pengolahan Hasil Perikanan Indonesia*, 21 (1): 42-53. <https://doi.org/10.17844/jphpi.v21i1.21261>
- Thomas, W. E.; Nilsson, L. M.; Forero, M.; Sokurenko, E. V.; Vogel, V., 2004: Shear-dependent stick and roll adhesion of type 1 fimbriated *Escherichia coli*. *Molecular Microbiology*, 53: 1545-1557. <https://doi.org/10.1111/j.1365-2958.2004.04226.x>
- Toma, M. M.; Pokrotieks, J., 2006: Prebiotics as functional food: Microbiological and Medical Aspects. *Acta Universitatis Latviensis*, 710: 117-129.
- Uraki, Y.; Koda, K., 2015: Utilization of wood cell wall components. *Journal of Wood Science*, 61 (5): 447-454. <https://doi.org/10.1007/s10086-015-1492-9>

Corresponding address:

SRI HIDAYATI

Lampung University, Agricultural Technology Department, Bandar Lampung, 35145, INDONESIA,
e-mail: srihidayati.unila@gmail.com